

Trigonometry (1/5): Introduction and Overview

Introduction to Engineering Mathematics


Prof. Joris Vankerschaver

Contents

- ① Angles and points on the unit circle
- ② Trigonometric functions as coordinates
- ③ Basic trigonometric identities

The unit circle

- The circle of radius 1 in the xy -plane, centered on the origin.
- Equation: $x^2 + y^2 = 1$
- Four quadrants: *I*, *II*, *III*, *IV*




Example

If $P(\sqrt{3}/2, y)$ is a point on the unit circle, find the value of y .

Angles and points on the unit circle

- Each point $P(x, y)$ defines an angle θ measured from the positive x -axis in counterclockwise direction.
- Angles measured in degrees or radians.
 - Value of θ in radians: length of arc subtended by θ (length of the red segment)




Converting between angles and radians

General formula to convert between degrees and radians:


$$\text{degrees} \begin{array}{c} \xrightarrow{\times \frac{\pi}{180}} \\ \xleftarrow{\times \frac{180}{\pi}} \end{array} \text{radians}$$

	Degrees	Radians
Full circle	360°	2π
Half circle	180°	π
Quarter circle	90°	$\pi/2$



Negative angles

Measured from the positive x -axis, in clockwise direction.




Adding 2π to an angle

- Point P is determined by the angle θ .
- P stays same when adding $\pm 2\pi$ to θ .

\Rightarrow All angles $\theta + 2k\pi$ with $k \in \mathbb{Z}$ give the same point P .


Principal angle: θ such that $-\pi < \theta \leq \pi$.



Finding the coordinates of a point

Given an angle θ , find the coordinates of $P(x, y)$.


- 1 $\theta = \pi/2$
- 2 $\theta = 3\pi$
- 3 $\theta = -\pi/2$



Finding the coordinates of a point

Slightly more involved case:

④ $\theta = \pi/4$




Important angles

Angle	x -coordinate	y -coordinate
0	1	0
$\pi/6$	$\sqrt{3}/2$	$1/2$
$\pi/4$	$\sqrt{2}/2$	$\sqrt{2}/2$
$\pi/3$	$1/2$	$\sqrt{3}/2$
$\pi/2$	0	1
π	-1	0
2π	1	0

Trigonometric functions as coordinates

Let θ be an angle with point $P(x, y)$.

Name	Notation	Definition
Cosine	$\cos \theta$	x
Sine	$\sin \theta$	y
Tangent	$\tan \theta$	$\frac{\sin \theta}{\cos \theta}$
Cotangent	$\cot \theta$	$\frac{\cos \theta}{\sin \theta}$
Cosecant	$\csc \theta$	$\frac{1}{\sin \theta}$
Secant	$\sec \theta$	$\frac{1}{\cos \theta}$




Example

Given that $\theta = \frac{\pi}{6}$, find the values of all 6 trigonometric functions.

Fundamental identity

- $P(x, y)$ is on the unit circle: $x^2 + y^2 = 1$
- Put $x = \cos \theta$ and $y = \sin \theta$ to obtain the **fundamental identity**:

$$\cos^2 \theta + \sin^2 \theta = 1$$



Aside: notation

Be very careful when you see $\sin^k \theta$.

- Positive exponent (**power**):

$$\sin^k \theta = (\sin \theta)^k.$$

- Negative exponent -1 (**inverse function**):

$$\sin^{-1} y = \arcsin y.$$

Fundamental identity: consequences

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

Example

Suppose $\cos \theta = -\frac{4}{5}$ and θ is in quadrant III. Find $\sin \theta$ and $\tan \theta$.

Periodicity of sine and cosine

- Sine and cosine are 2π -periodic:

$$\sin(\theta \pm 2\pi) = \sin \theta$$

$$\cos(\theta \pm 2\pi) = \cos \theta$$

- The tangent is π -periodic:

$$\tan(\theta \pm \pi) = \tan \theta$$

Example: Compute $\tan\left(\frac{8093\pi}{4}\right)$

