Coordinate Geometry Introduction to Engineering Mathematics


Prof. Joris Vankerschaver

Contents

- 1 Coordinate geometry
- 2 Locus of points
- 3 Equations of circles
- **4** Equations of lines
- 6 Exercise

What is coordinate geometry?

Studying geometry through coordinate calculations.






Example: distance between two points

Distance between P and Q:


$$d(P,Q)=\sqrt{(x_2-x_1)^2+(y_2-y_1)^2}$$



Example: midpoint between two points

Coordinates of midpoint between \boldsymbol{P} and $\boldsymbol{Q}:$

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right).$$




Locus of points

"Locus" = Set of points satisfying some condition.

- **Circle**: All points at given distance from a fixed center.
- **Ellipse**: All points for which the sum of distances to two fixed points (focal points) is constant
- **Parabola**: All points that are at equal distance from a fixed point and a given line (directrix)





Find the locus of points for which the distance to the x-axis is equal to the distance to the point (0, 1).


Circles

Locus of points P(x,y) at distance R from center C(h,k). We have d(P,C) = R so that

$$\sqrt{(x-h)^2+(y-k)^2}=R,$$

and by squaring

$$(x-h)^2 + (y-k)^2 = R^2$$



Find the equation of the circle that has the points $\left(1,1\right)$ and $\left(7,9\right)$ as end points of a diameter.

Find the center and radius of the circle given by $x^2 + y^2 - 6x + 2y + 8 = 0.$


Lines

Line not parallel to the y-axis:

$$y = mx + q$$

with

- *m*: the **slope**
- q: the intercept




Lines

Line parallel to the y-axis:

$$x = a$$

with

• *a*: where the line intersects the *x*-axis




Finding the slope of a line

Take

$$\begin{array}{l} \bullet \ \Delta x = x_2 - x_1 \\ \bullet \ \Delta y = y_2 - y_1 \\ \end{array} \\ \mbox{Then} \end{array}$$

$$m = \frac{\Delta y}{\Delta x} = \frac{y_2 - y_1}{x_2 - x_1}$$

" Δx steps to the right, Δy steps up/down."



Find the equation for the line through (1,5) and (2,7).

Properties

Equation for the line through (x_0, y_0) with slope m:

$$y-y_0=m(x-x_0)$$


Equation for the line through the points (x_1, y_1) and (x_2, y_2) :

$$y-y_1=\frac{y_2-y_1}{x_2-x_1}(x-x_1).$$

Parallel/perpendicular lines

Two lines are ...

- parallel if their slopes are the same: $m_1 = m_2$
- perpendicular if their slopes satisfy: $\left| m_1 m_2 = -1 \right|$



In general, the angle θ between two lines is determined by

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|.$$


Given two lines $L_1:x+2y-3=0$ and $L_2:kx+y-5=0,$ for which value of k are L_1 and $L_2\ldots$

- 1 Parallel?
- Perpendicular?
- **3** At an angle of 45° ?

Distance of a point to a line

Distance between point $P(x_0, y_0)$ and line L: ax + by + c = 0:

$$D = \frac{|ax_0 + by_0 + c|}{\sqrt{a^2 + b^2}}.$$



Different representations of lines


- Slope/intercept:
 - y = mx + q (not parallel to y-axis)
 - x = a (parallel)
- **2** Linear representation: ax + by + c = 0
- **3** Polar representation:
 - For line through the origin: $\tan \theta = m$
 - For line not through the origin:

$$r = \frac{q}{\sin \theta - m \cos \theta}$$

Exercise

Find the equation of the common tangent line between two touching circles given by

$$\begin{split} C_1 &: x^2 + y^2 - 6x - 12y + 37 = 0 \\ C_2 &: x^2 + y^2 - 6y + 7 = 0. \end{split}$$



Source: xkcd 2735